

# Bladder cancer risk in painters: A meta-analysis - Abstract

Contributed by Administrator  
Tuesday, 10 August 2010

Section of IARC Monographs, 150 cours Albert Thomas, 69372 Lyon Cedex 08, France.

guhan@iarc.fr

The International Agency for Research on Cancer has classified occupational exposure as a painter as 'carcinogenic to humans', largely based on increased risks of bladder and lung cancer. A meta-analysis, including more than 2900 incident cases or deaths from bladder cancer among painters reported in 41 cohort (n=2), record linkage (n=9) and case-control (n=30) studies, was conducted to quantitatively compare the results of the different study designs and the potential confounding effect of smoking as well as other occupational exposures. The summary relative risk (meta-RR, random effects) for bladder cancer in painters was 1.25 (95% CI 1.16 to 1.34; 41 studies) overall and 1.28 (95% CI 1.15 to 1.43; 27 studies) when including only smoking adjusted risk estimates. The elevated risk persisted when restricted to studies that adjusted for other occupational exposures (meta-RR 1.27; 95% CI 0.99 to 1.63; 4 studies). The results remained robust when stratified by study design, gender and study location. Furthermore, exposure-response analyses suggested that the risk increased with duration of employment. There was no evidence of publication bias. Taken together, these results support the conclusion that occupational exposures in painters are causally associated with the risk of bladder cancer.

Written by:

Guha N, Steenland NK, Merletti F, Altieri A, Coglianò V, Straif K.  
&#160;&#160;Are you the author?

Reference: *Occup Environ Med.* 2010 Aug;67(8):568-73.

doi: 10.1136/oem.2009.051565

PubMed Abstract

PMID: 20647380

UroToday.com Bladder Cancer Section

